

PRODUCT INFORMATION

Clone ID	DMC688
Target	ASGR1
Synonyms	ASGPR; ASGPR1; CLEC4H1; HL-1
Host Species	Rabbit
Description	Anti-ASGR1 antibody(DMC688); IgG1 Chimeric mAb
Delivery	In Stock
Uniprot ID	P07306; Q6FGQ5
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a subunit of the asialoglycoprotein receptor. This receptor is a transmembrane protein that plays a critical role in serum glycoprotein homeostasis by mediating the endocytosis and lysosomal degradation of glycoproteins with exposed terminal galactose or N-acetylgalactosamine residues. The asialoglycoprotein receptor may facilitate hepatic infection by multiple viruses including hepatitis B; and is also a target for liver-specific drug delivery. The asialoglycoprotein receptor is a hetero-oligomeric protein composed of major and minor subunits; which are encoded by different genes. The protein encoded by this gene is the more abundant major subunit. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq; Jan 2011]
Usage	Research use only






Figure 1. Flow cytometry analysis with Anti-ASGR1(DMC688) on Expi293 cells transfected with human ASGR1(Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human ASGR1 mAb(DMC100688).

(A) DMC100688 does not bind to 293T cells that do not express ASGR1.

(B) A clear peak shift of DMC100688 was seen compared to the control when incubated with ASGR1-expressing HepG2 cells, indicating strong binding of DMC100688 to ASGR1. Antibodies were incubated at 5 μ g/mL.

