

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM110
<b>Target</b>	B7-1
<b>Synonyms</b>	CD80;B7;B7-1;B7.1;BB1;CD28LG;CD28LG1;LAB7
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-B7-1 antibody(DM110); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P33681
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a membrane receptor that is activated by the binding of CD28 or CTLA-4. The activated protein induces T-cell proliferation and cytokine production. This protein can act as a receptor for adenovirus subgroup B and may play a role in lupus neuropathy.
<b>Usage</b>	Research use only






Figure 1. ELISA plate pre-coated by 2  $\mu\text{g/ml}$  (100  $\mu\text{l/well}$ ) Human B7-1 protein, hFc tagged protein PME100473 can bind Rabbit anti-B7-1 monoclonal antibody (clone: DM110) in a linear range of 0.32-100 ng/ml.

Figure 2. Flow cytometry analysis with Anti-B7-1 (DM110) on Expi293 cells transfected with human B7-1(Blue histogram) or Expi293 transfected with irrelevant protein(Red histogram).




Figure 3. Flow cytometry analysis of antigen binding of rabbit anti-human B7-1 mAb(DME100110).

(A) DME100110 does not bind to CHO-S cells that do not express B7-1.

(B) A clear peak shift of DME100110 was seen compared to the control when incubated with B7-1-expressing Raji cells, indicating strong binding of DME100110 to B7-1. Antibodies were incubated at 5  $\mu\text{g/mL}$ .

