

PRODUCT INFORMATION

Clone ID	8B5
Target	CD22
Synonyms	SIGLEC2; SIGLEC-2
Host Species	Rabbit
Description	Anti-CD22 antibody(8B5), Rabbit mAb
Delivery	In Stock
Uniprot ID	P20273
lgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	WB; Flow Cyt
Recommended Dilutions	WB 1:1000; Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	CD22 (CD22 Molecule) is a Protein Coding gene. Diseases associated with CD22 include Refractory Hematologic Cancer and Hairy Cell Leukemia. Among its related pathways are Downstream signaling events of B Cell Receptor (BCR) and Hematopoietic cell lineage. Gene Ontology (GO) annotations related to this gene include carbohydrate binding. An important paralog of this gene is SIGLEC2.
Usage	Research use only
Conjugate	Unconjugated

Email: info@dimabio.com Website: www.dimabio.com





130 kDa -	-
100 kDa -	
70 kDa -	
55 kDa -	
35 kDa -	

25 kDa -

Figure 1.Anti-CD22 antibody (SKU# DME100012) at 1/1000 dilution

Lane : Raji (human Burkitt's lymphoma B lymphocyte), whole cell lysate

Secondary : Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 60 kDa Observed band size: 130 kDa




Figure 2. Flow cytometry analysis of antigen binding of anti-human CD22 mAb(DME100012).

(A) DME100012 does not bind to Jurkat cells that do not express CD22
(B) A clear peak shift of DME100012 was seen compared to the control when incubated with CD22-expressing Raji cells, indicating strong binding of DME100012 to CD22.
Antibodies were incubated at 10 ug/mL.

Email: info@dimabio.com Website: www.dimabio.com

