

**PRODUCT INFORMATION**

<b>Common Name</b>	HuMax-CD38
<b>Synonyms</b>	CD38;T10;ADPRC1
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Homo sapiens
<b>IgG type</b>	IgG1
<b>Reactivity</b>	Human
<b>Target</b>	CD38
<b>Uniprot ID</b>	P28907
<b>Description</b>	Anti-CD38 (daratumumab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only



### Anti-CD38 (daratumumab biosimilar) mAb ELISA

0.2  $\mu$ g Human CD38, hFc-His Tagged protein per well




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ L/well) Human CD38, hFc-His tagged protein (PME100004) can bind Anti-CD38 Neutralizing antibody in a linear range of 0.98-64.14 ng/ml. In order to specifically detect BME100004, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. HEK293 cell line transfected with irrelevant protein (**B**) and human CD38 (**A**) were surface stained with anti-CD38 neutralizing antibody 1 $\mu$ g/ml (daratumumab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.

