

PRODUCT INFORMATION

Clone ID	DM191
Target	CD70
Synonyms	CD70;CD27LG;TNFSF7;TNFSF7G;CD27L
Host Species	Rabbit
Description	Anti-CD70 antibody(DM191); Rabbit mAb
Delivery	In Stock
Uniprot ID	P32970
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27:CD27. It is a surface antigen on activated; but not on resting; T and B lymphocytes. It induces proliferation of costimulated T cells; enhances the generation of cytolytic T cells; and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation; cytotoxic function of natural killer cells; and immunoglobulin synthesis.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Flow cytometry analysis with Anti-CD70 (DM191) on Expi293 cells transfected with human CD70 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

