

PRODUCT INFORMATION

Common Name	CD74-DOX (ADC),hLL1,hLL1-DOX (ADC),MEDI-115
Synonyms	DHLAG
Applications	ELISA, Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Host Species	Humanized
IgG type	IgG1(K97R)
Reactivity	Human
Target	CD74
Uniprot ID	P04233
Description	Anti-CD74(milatumzumab biosimilar) mAb
Delivery	In Stock
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only




Figure 1. Flow cytometry analysis with 15 μ g/mL Anti-CD74(milatuzumab biosimilar) mAb (BME100170) on Expi293 cells transfected with Human CD74 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Anti-CD74(milatuzumab biosimilar) mAb ELISA

0.2 μ g of Human CD74, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CD74 (73-296) Protein, hFc Tag(PME100642) can bind Anti-CD74(milatuzumab biosimilar) mAb(BME100170) in a linear range of 0.64-16 ng/mL. In order to specifically detect BME100170, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 3. Flow cytometry analysis of antigen binding of anti-human CD74 mAb(BME100170).

(A) BME100170 does not bind to 293T cells that do not express CD74.

(B) A clear peak shift of BME100170 was seen compared to the control when incubated with CD74-expressing Raji cells, indicating strong binding of BME100170 to CD74. Antibodies were incubated at 5 μ g/mL.

