Anti-CD79B antibody(52A11); IgG1 Chimeric mAb

Cat. No. DMC101033



PRODUCT INFORMATION

Clone ID 52A11 CD79B **Target**

Synonyms AGM6;B29;IGB

Host Species Rabbit

Description Anti-CD79B antibody(52A11); IgG1 Chimeric mAb

Delivery In Stock **Uniprot ID** P40259

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human

Applications Flow Cyt; WB

Recommended

Background

Flow Cyt 1:100; WB 1:1000 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-

cell antigen receptor. This gene encodes the Igbeta protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described. [provided

> Email: info@dimabio.com Website: www.dimabio.com

by RefSeq, Jul 2008]

Usage Research use only

Conjugate Unconjugated






Figure 1. Flow cytometry analysis with Anti-CD79B (52A11) mAb on Expi293 cells transfected with human CD79B (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).





Figure 2. Flow cytometry analysis of antigen binding of anti-human CD79B mAb(DMC101033). (A) DMC101033 does not bind to 293T cells that do not express CD79B. (B) A clear peak shift of DMC101033 was seen compared to the control when incubated with CD79B-expressing Raji cells, indicating strong binding of DMC101033 to CD79B. Antibodies were incubated at 5 μ g/mL.



Email: info@dimabio.com Website: www.dimabio.com



Anti-CD79B antibody(52A11); IgG1 Chimeric mAb

Cat. No. DMC101033



250 kDa-130 kDa-100 kDa-70 kDa-55 kDa-25 kDa-

Figure 3.Anti-CD79B antibody (SKU# DMC101033) at 1/1000 dilution

Lane: RAJI (human Burkitt's lymphoma B lymphocyte), whole cell lysate

Email: info@dimabio.com Website: www.dimabio.com

Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 26 kDa Observed band size: 24 kDa

