

PRODUCT INFORMATION

BMS-936564, MDX-1338 **Common Name**

Conjugate Unconjugated

CD184;D2S201E;FB22;HM89;HSY3RR;LAP-3;LAP3;LCR1;LESTR;NPY3R;NPYR;NPYRL;NPYY3R;WHIM;WHIMS Synonyms

Applications Flow Cyt Recommended Dilutions Flow Cyt 1:100

 $Lyophilized from sterile \ PBS, pH\ 7.4.\ Normally\ 5\%-8\%\ trehalose\ is\ added\ as\ protectants\ before\ lyophilization.$ Formulation & Reconstitution

Host Species IgG type IgG4 Reactivity Human Target CXCR4

Description Anti-CXCR4(ulocuplumab biosimilar) mAb

P61073

Delivery In Stock

Uniprot ID

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. Storage & Shipping

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.

Usage Research use only

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. **DIMA Disclaimer**




Figure 1. Flow cytometry analysis with 15 μ g/mL Anti-CXCR4 (ulocuplumab biosimilar) mAb (BME100101) on Expi293 cells transfected with Human CXCR4 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com









Figure 2. Flow cytometry analysis of antigen binding of anti-human CXCR4 mAb(BME100101). (A) BME100101 does not bind to hepG2 cells that do not express CXCR4. (B) A clear peak shift of BME100101 was seen compared to the control when incubated with CXCR4-expressing Hela cells, indicating strong binding of BME100101 to CXCR4. Antibodies were incubated at 5 μ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

