

PRODUCT INFORMATION

BMS-936564, MDX-1338 **Common Name**

CD184;D2S201E;FB22;HM89;HSY3RR;LAP-3;LAP3;LCR1;LESTR;NPY3R;NPYR;NPYRL;NPYY3R;WHIM;WHIMSSynonyms

Applications Flow Cyt Recommended Dilutions Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Host Species Homo sapiens IgG type lgG4 Reactivity Human CXCR4 Target Uniprot ID P61073

Anti-CXCR4(ulocuplumab biosimilar) mAb Description

Delivery In Stock

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature $\frac{1}{2}$ Storage & Shipping

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals. Background

Usage Research use only




Figure 1. Flow cytometry analysis with 15 μ g/mL Anti-CXCR4 (ulocuplumab biosimilar) mAb (BME100101) on Expi293 cells transfected with Human CXCR4 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com









Figure 2. Flow cytometry analysis of antigen binding of anti-human CXCR4 mAb(BME100101). (A) BME100101 does not bind to hepG2 cells that do not express CXCR4. (B) A clear peak shift of BME100101 was seen compared to the control when incubated with CXCR4-expressing Hela cells, indicating strong binding of BME100101 to CXCR4. Antibodies were incubated at 5 μ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

