

PRODUCT INFORMATION

Clone ID	BM301
Target	DDK
Synonyms	ddddk; dykddddk; flag
Host Species	Mouse
Description	Anti-DDK antibody(BM301); Mouse IgG1 mAb
Delivery	In Stock
Uniprot ID	N/A
IgG type	IgG1
Clonality	Monoclonal
Reactivity	Flag is widely used as an additional tag in expression vectors. The tag can be fused with the target protein at the N-terminal and C-terminal, and can be detected.
Applications	ELISA, WB
Recommended Dilutions	ELISA 1:5000-10000, WB 1:1000
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
Background	Flag is widely used as an additional tag in expression vectors. The tag can be fused with the target protein at the N-terminal and C-terminal, and can be detected.
Usage	Research use only
Conjugate	Unconjugated

