

PRODUCT INFORMATION

Clone ID	DMC460
Target	FGFR4
Synonyms	CD334; JTK2; TKF
Host Species	Rabbit
Description	Anti-FGFR4 antibody(DMC460); IgG1 Chimeric mAb
Delivery	In Stock
Uniprot ID	P22455
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a tyrosine kinase and cell surface receptor for fibroblast growth factors. The encoded protein is involved in the regulation of several pathways; including cell proliferation; cell differentiation; cell migration; lipid metabolism; bile acid biosynthesis; vitamin D metabolism; glucose uptake; and phosphate homeostasis. This protein consists of an extracellular region; composed of three immunoglobulin-like domains; a single hydrophobic membrane-spanning segment; and a cytoplasmic tyrosine kinase domain. The extracellular portion interacts with fibroblast growth factors; setting in motion a cascade of downstream signals; ultimately influencing mitogenesis and differentiation. [provided by RefSeq; Aug 2017]
Usage	Research use only






Figure 1. Flow cytometry analysis with Anti-FGFR4 (DMC460) on Expi293 cells transfected with human FGFR4 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human FGFR4 mAb(DMC100460).

(A) DMC100460 does not bind to SiHa cells that do not express FGFR4.

(B) A clear peak shift of DMC100460 was seen compared to the control when incubated with FGFR4-expressing Huh7 cells, indicating strong binding of DMC100460 to FGFR4. Antibodies were incubated at 5 μ g/mL.

