

## **PRODUCT INFORMATION**

Clone ID 19A5 FLT3 **Target** 

**Synonyms** Flt-3;Flk-2;STK-1;CD135;FLK2;FLT-3

**Host Species** Rabbit

Description Anti-FLT3 antibody(19A5), IgG1 Chimeric mAb

**Delivery** In Stock **Uniprot ID** P36888

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

**Background** 

Flow Cyt 1/100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Flt-3(Receptor-type tyrosine-protein kinase FLT3) is also known as FLK-2(Fetal liver kinase-2), STK-1(Stem cell tyrosine kinase 1), CD135. FLT3 is a cytokine receptor which belongs to the receptor tyrosine kinase class III. Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine FLT3LG and regulates differentiation, proliferation and survival of hematopoietic progenitor cells and of dendritic cells. Promotes phosphorylation of SHC1 and AKT1, and

activation of the downstream effector MTOR. Promotes activation of RAS signaling and

phosphorylation of downstream kinases, including MAPK1/ERK2 and/or MAPK3/ERK1. Mutations that cause constitutive kinase activity promote cell proliferation and resistance to apoptosis via the activation of multiple signaling pathways. Research use only All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application.

> Email: info@dimabio.com Website: www.dimabio.com

Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing Usage

all patent application to ensure no IP

infringement.

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)









Figure 1. Flow cytometry analysis with 1µg/mL Anti-FLT3 (19A5) mAb on THP-1 cells.



Email: info@dimabio.com Website: www.dimabio.com

