

**PRODUCT INFORMATION**

<b>Clone ID</b>	55A1
<b>Target</b>	HER3
<b>Synonyms</b>	ERBB3; FERLK; LCCS2; VSCN1; ErbB-3; c-erbB3; erbB3-S; MDA-BF-1; c-erbB-3; p180-ErbB3; p45-sErbB3; p85-sErbB3
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-Her3 antibody(55A1), Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P21860
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	IHC; Flow Cyt
<b>Recommended Dilutions</b>	IHC 1:100; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been characterized. One isoform lacks the intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized. [provided by RefSeq, Jul 2008]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Flow cytometry analysis of antigen binding of anti-human Her3 mAb(DME100580 ).

(A) DME100580 does not bind to 293T cells that do not express Her3

(B) A clear peak shift of DME100580 was seen compared to the control when incubated with Her3-expressing Huh7 cells, indicating strong binding of DME100580 to Her3.

Antibodies were incubated at 10  $\mu$ g/mL.

