



**PRODUCT INFORMATION**

<b>Clone ID</b>	DM132
<b>Target</b>	HVEM
<b>Synonyms</b>	ATAR; CD270; HVEA; HVEM; LIGHTR; TR2
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-HVEM antibody(DM132); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q92956
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of the TNF (tumor necrosis factor) receptor superfamily. The encoded protein functions in signal transduction pathways that activate inflammatory and inhibitory T-cell immune response. It binds herpes simplex virus (HSV) viral envelope glycoprotein D (gD); mediating its entry into cells. Alternative splicing results in multiple transcript variants.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated





**Figure 1.** ELISA plate pre-coated by 1  $\mu\text{g/ml}$  (100  $\mu\text{l/well}$ ) Human HVEM protein, His tagged protein ([getskuurl sku="PME100273"]) can bind Rabbit anti-HVEM monoclonal antibody(clone: **DM132**) in a linear range of 0. 1-12 ng/ml.



**Figure 2.** Flow cytometry analysis with Anti-HVEM (**DM132**) on Expi293 cells transfected with human HVEM(Blue histogram) or Expi293 transfected with irrelevant protein(Red histogram).

