

PRODUCT INFORMATION

Common Name MDX-1333, MEDI-546, anifrolumab-fnia

Conjugate Unconjugated

Synonyms IFN-R-1;CRF2-1;IFNAR

Applications ELISA; Flow Cyt

Recommended

Dilutions

ELISA 1:5000-10000; Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

Host Species Homo sapiens

IgG type lgG1 Reactivity Human **Target** IFNAR1 **Uniprot ID** P17181

Anti-IFNAR1(anifrolumab biosimilar) mAb **Description**

Delivery In Stock

Storage & Shipping

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in

Background therapeutic or diagnostic procedures for humans

or animals.


Usage Research use only

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

DIMA Disclaimer

actively scrutinizing all patent application to ensure no IP infringement.









Anti-IFNAR1 (anifrolumab biosimilar) mAb ELISA

0.1 μg of Human IFNAR1, His tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/mL (100 μ L/well) Human IFNAR1 Protein, His Tag ([getskuurl sku="PME100958"]) can bind Anti-IFNAR1 Neutralizing antibody (BME100117) in a linear range of 0.61–39.06 ng/mL.

Anti-IFNAR1 (anifrolumab biosimilar) mAb ELISA

0.2 µg of Human IFNAR1, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human IFNAR1 Protein, hFc Tag (PME100773) can bind Anti-IFNAR1 Neutralizing antibody (BME100117) in a linear range of 0.73–187.50 ng/mL. In order to specifically detect BME100117, mouse anti-human Fab-specific antibody was used as detection antibody.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China) Email: info@dimabio.com Website: www.dimabio.com





Anti-IFNAR1 (anifrolumab biosimilar) mAb ELISA

0.2 μg of Human IFNAR1, mFc tagged protein per well




Figure 3. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human IFNAR1 Protein, mFc Tag ([getskuurl sku="PME100737"]) can bind Anti-IFNAR1 Neutralizing antibody (BME100117) in a linear range of 1.22–312.50 ng/mL.




Figure 4. Flow cytometry analysis with 1 μ g/mL Anti-IFNAR1 (anifrolumab biosimilar) mAb (BME100117) on Expi293 cells transfected with Human IFNAR1 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

