Cat. No. BME100114



PRODUCT INFORMATION

Common Name ILV-094

Conjugate Unconjugated

Synonyms IL-22; Cytokine Zcyto18; IL-TIF

Applications ELISA; Flow Cyt

Recommended

Dilutions

ELISA 1:5000-10000; Flow Cyt 1:100

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution.

Host Species Homo sapiens

IgG type lgG1 Reactivity Human **Target** IL22 **Uniprot ID** Q9GZX6

Anti-IL22(fezakinumab biosimilar) mAb **Description**

Delivery In Stock

> Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in

Background therapeutic or diagnostic procedures for humans

or animals.

Usage Research use only

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

> > Email: info@dimabio.com Website: www.dimabio.com

DIMA Disclaimer

actively scrutinizing all patent application to ensure no IP infringement.





Anti-IL22 (fezakinumab biosimilar) mAb ELISA

0.2 μg of Human IL22, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human IL22 Protein, hFc Tag (PME100651) can bind Anti-IL22 Neutralizing antibody (BME100114) in a linear range of 2.44–625.00 ng/mL. In order to specifically detect BME100114, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis under cell membrane permeable condition with 1 μ g/mL Anti-IL22 (fezakinumab biosimilar) mAb (BME100114) on Expi293 cells transfected with Human IL22 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com