Cat. No. BME100021



PRODUCT INFORMATION

Common Name MORAb-009 Conjugate Unconjugated

Synonyms MSLN; Mesothelin; MPF

Applications ELISA; Flow Cyt

Recommended **Dilutions**

ELISA 1:5000-10000; Flow Cyt 1:100

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution.

Host Species Chimeric IgG type lgG1 Reactivity Human **Target** MSLN **Uniprot ID** Q13421

Anti-Mesothelin (amatuximab biosimilar) mAb **Description**

Delivery In Stock

> Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in

Background therapeutic or diagnostic procedures for humans

or animals.

Usage Research use only

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

DIMA Disclaimer

actively scrutinizing all patent application to ensure no IP infringement.

Email: info@dimabio.com Website: www.dimabio.com





Anti-Mesothelin (amatuximab biosimilar) mAb ELISA

0.2 µg of Mesothelin(296-580), mFc-His Tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human Mesothelin, mFc-His tagged protein ([getskuurl sku="PME100031"]) can bind Anti-Mesothelin Neutralizing antibody in a linear range of 3.2-80.0 ng/ml.




Figure 2. HEK293 cell line transfected with irrelevant protein (B) and human Mesothelin (A) were surface stained with anti-Mesothelin neutralizing antibody $1\mu g/ml$ (amatuximab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.

Email: info@dimabio.com Website: www.dimabio.com

