


**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Clone ID</b>                         | DM168  |
| <b>Target</b>                           | MICB   |
| <b>Synonyms</b>                         | MIC-B; PERB11.2  |
| <b>Host Species</b>                     | Rabbit   |
| <b>Description</b>                      | Anti-MICB antibody(DM168); Rabbit mAb  |
| <b>Delivery</b>                         | In Stock   |
| <b>Uniprot ID</b>                       | Q29980   |
| <b>IgG type</b>                         | Rabbit IgG   |
| <b>Clonality</b>                        | Monoclonal   |
| <b>Reactivity</b>                       | Human  |
| <b>Applications</b>                     | ELISA; Flow Cyt  |
| <b>Recommended Dilutions</b>            | ELISA 1:5000-10000; Flow Cyt 1:100   |
| <b>Purification</b>                     | Purified from cell culture supernatant by affinity chromatography  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.   |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Background</b>                       | This gene encodes a heavily glycosylated protein which is a ligand for the NKG2D type II receptor. Binding of the ligand activates the cytolytic response of natural killer (NK) cells; CD8 alphabeta T cells; and gammadelta T cells which express the receptor. This protein is stress-induced and is similar to MHC class I molecules; however; it does not associate with beta-2-microglobulin or bind peptides. Alternative splicing results in multiple transcript variants. |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Unconjugated   |





**Figure 1.** MICB protein is highly expressed on the surface of Expi293 cell membrane. Flow cytometry analysis with Anti-MICB (DM168) on Expi293 cells transfected with human MICB (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram), and Isotype antibody on Expi293 transfected with irrelevant protein (Orange histogram).

DIMABIO CONFIDENTIAL

