

## **PRODUCT INFORMATION**

Clone ID **DMC267 Target** TNFSF11

CD254; hRANKL2; ODF; OPGL; OPTB2; RANKL; **Synonyms** 

sOdf; TNLG6B; TRANCE

**Host Species** 

Anti-TNFSF11 antibody(DMC267); IgG1 Chimeric **Description** 

mAb In Stock

014788 IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Background

Delivery

**Uniprot ID** 

Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the tumor necrosis factor (TNF) cytokine family which is a ligand for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. This protein was shown to be a dentritic cell survival factor and is involved in the regulation of T cell-dependent immune response. T cell activation was reported to induce expression of

this gene and lead to an increase of osteoclastogenesis and bone loss. This protein

was shown to activate antiapoptotic kinase AKT:PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor-associated factor (TRAF) 6; which indicated this protein may have a role in the regulation of cell apoptosis. Targeted disruption of the related gene in mice led to severe osteopetrosis and a lack of osteoclasts. The deficient mice exhibited defects in early differentiation of T and B lymphocytes; and failed to form lobulo-alveolar mammary

structures during pregnancy. Two alternatively spliced transcript variants have been found.


Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





**Figure 1.** Flow cytometry analysis with Anti-TNFSF11 (DMC267) on Expi293 cells transfected with human TNFSF11 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

