

## **PRODUCT INFORMATION**

IMMU-132, hRS7-SN-38, hRS7-SN-38-ADC, hRS7-**Common Name** 

[CL-SN-38], hrS7-SN-38

**Synonyms** TACSTD2;GA733-1;M1S1;TROP2

**Applications** ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions** 

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

**Host Species** Humanized

IgG type lgG1 Reactivity Human **Target** Trop2 **Uniprot ID** P09758

**Description** Anti-Trop2 (sacituzumab biosimilar) mAb

**Delivery** In Stock

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

Email: info@dimabio.com Website: www.dimabio.com

temperature.

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans **Background** 

or animals.

Usage Research use only

Conjugate Unconjugated





## Anti-Trop2 (sacituzumab govitecan biosimilar) mAb ELISA

0.2 µg of Trop2, mFc-His Tagged protein per well




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human Trop2, mFc-His tagged protein PME100501 can bind Anti-Trop2 Neutralizing antibody in a linear range of 0.13-16.0 ng/ml.




Figure 2. Flow cytometry analysis of antigen binding of anti-human Trop2 mAb(BME100023). (A) BME100023 does not bind to 293T cells that do not express Trop2. (B) A clear peak shift of BME100023 was seen compared to the control when incubated with Trop2-expressing A431 cells, indicating strong binding of BME100023 to Trop2. Antibodies were incubated at 2  $\mu$ g/mL.

Email: info@dimabio.com Website: www.dimabio.com








Figure 3. HEK293 cell line transfected with irrelevant protein (B) and human Trop2 protein (A) were surface stained with anti-Trop2 neutralizing antibody  $1\mu g/ml$  (sacituzumab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.



