


**PRODUCT INFORMATION**

<b>Clone ID</b>	DM185
<b>Target</b>	VEGFR2
<b>Synonyms</b>	CD309; FLK1; VEGFR; VEGFR2
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-VEGFR2 antibody(DM185); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P35968
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Storage &amp; Shipping</b>	
<b>Background</b>	Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor; known as kinase insert domain receptor; is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation; survival; migration; tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors; including Rab GTPase; P2Y purine nucleotide receptor; integrin alphaVbeta3; T-cell protein tyrosine phosphatase; etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.





**Figure 1.** Flow cytometry analysis with Anti-VEGFR2 (DM185) on Expi293 cells transfected with human VEGFR2 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

