

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Clone ID</b>                         | DM138   |
| <b>Target</b>                           | 5T4   |
| <b>Synonyms</b>                         | 5T4   |
| <b>Host Species</b>                     | Rabbit  |
| <b>Description</b>                      | Biotinylated Anti-5T4 antibody(DM138); Rabbit mAb   |
| <b>Delivery</b>                         | 2-3 weeks   |
| <b>Uniprot ID</b>                       | Q13641  |
| <b>IgG type</b>                         | Rabbit IgG  |
| <b>Clonality</b>                        | Monoclonal  |
| <b>Reactivity</b>                       | Human   |
| <b>Applications</b>                     | ELISA; Flow Cyt   |
| <b>Recommended Dilutions</b>            | ELISA 1:5000-10000; Flow Cyt 1:100  |
| <b>Purification</b>                     | Purified from cell culture supernatant by affinity chromatography   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.   |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | This gene encodes a leucine-rich transmembrane glycoprotein that may be involved in cell adhesion. The encoded protein is an oncofetal antigen that is specific to trophoblast cells. In adults this protein is highly expressed in many tumor cells and is associated with poor clinical outcome in numerous cancers. Alternate splicing in the 5' UTR results in multiple transcript variants that encode the same protein. |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Biotinylated  |

