Cat. No. DMC100446B



PRODUCT INFORMATION

Clone ID **DMC446** CD117 **Target**

Synonyms C-Kit; CD117; MASTC; PBT; SCFR; KIT

Host Species

Biotinylated Anti-CD117 antibody(DMC446); IgG1 Description

Chimeric mAb

Delivery 2-3 weeks **Uniprot ID** P10721

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Background

Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a receptor tyrosine kinase. This gene was initially identified as a homolog of the feline sarcoma viral oncogene v-kit and is

often referred to as proto-oncogene c-Kit. The canonical form of this glycosylated transmembrane protein has an N-terminal extracellular region with five immunoglobulin-like domains; a transmembrane region; and an intracellular tyrosine kinase domain at the C terminus. Upon activation by its cytokine ligand; stem cell factor (SCF); this protein phosphorylates multiple intracellular proteins that play a role in the proliferation; differentiation; migrative plays and therefore the province of many cell types and therefore plays are

apoptosis of many cell types and thereby plays an important role in hematopoiesis; stem cell maintenance; gametogenesis; melanogenesis; and in mast cell development; migration and function. This protein can be a membrane-bound or soluble protein. Mutations in this gene are associated with gastrointestinal stromal tumors; mast cell disease; acute myelogenous leukemia;

and piebaldism. Multiple transcript variants encoding different isoforms have been found for

> Email: info@dimabio.com Website: www.dimabio.com

this gene.

Usage Research use only

Biotinylated Conjugate

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)