

## PRODUCT INFORMATION

<b>Clone ID</b>	DMC446
<b>Target</b>	CD117
<b>Synonyms</b>	C-Kit; CD117; MASTC; PBT; SCFR; KIT
<b>Host Species</b>	Rabbit
<b>Description</b>	Biotinylated Anti-CD117 antibody(DMC446); IgG1 Chimeric mAb
<b>Delivery</b>	2-3 weeks
<b>Uniprot ID</b>	P10721
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a receptor tyrosine kinase. This gene was initially identified as a homolog of the feline sarcoma viral oncogene v-kit and is often referred to as proto-oncogene c-Kit. The canonical form of this glycosylated transmembrane protein has an N-terminal extracellular region with five immunoglobulin-like domains; a transmembrane region; and an intracellular tyrosine kinase domain at the C-terminus. Upon activation by its cytokine ligand; stem cell factor (SCF); this protein phosphorylates multiple intracellular proteins that play a role in the proliferation; differentiation; migration and apoptosis of many cell types and thereby plays an important role in hematopoiesis; stem cell maintenance; gametogenesis; melanogenesis; and in mast cell development; migration and function. This protein can be a membrane-bound or soluble protein. Mutations in this gene are associated with gastrointestinal stromal tumors; mast cell disease; acute myelogenous leukemia; and piebaldism. Multiple transcript variants encoding different isoforms have been found for this gene.
<b>Usage</b>	Research use only

