

## PRODUCT INFORMATION

DMC269 Clone ID CD44 Target

CD44;CDW44;CSPG8;ECMR-III;HCELL;HUTCH-I;IN;LHR;MC56;MDU2;MDU3;MIC4;Pgp1;Epican Synonyms

**Host Species** 

Description Biotinylated Anti-CD44 antibody(DMC269); IgG1 Chimeric mAb

Delivery 2-3 weeks P16070 **Uniprot ID** 

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human Applications Flow Cyt Recommended Dilutions Flow Cyt 1:100

Purified from cell culture supernatant by affinity chromatography Purification

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell interactions; cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands; such as osteopontin; collagens; and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation; recirculation and homing; hematopoiesis; and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms; however; the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein; and may be related to tumor metastasis.

Email: info@dimabio.com Website: www.dimabio.com

Usage Conjugate Biotinylated

**Background** 

