

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM203
<b>Target</b>	CD99
<b>Synonyms</b>	HBA71; MIC2; MIC2X; MIC2Y; MSK5X
<b>Host Species</b>	Rabbit
<b>Description</b>	Biotinylated Anti-CD99 antibody(DM203); Rabbit mAb
<b>Delivery</b>	2-3 weeks
<b>Uniprot ID</b>	P14209
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte migration; T-cell adhesion; ganglioside GM1 and transmembrane protein transport; and T-cell death by a caspase-independent pathway. In addition; the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. There is a related pseudogene located immediately adjacent to this locus. [provided by RefSeq; Mar 2016]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Biotinylated

