

## **PRODUCT INFORMATION**

Clone ID **DMC366 Target CFB** 

AHUS4; ARMD14; BF; BFD; CFAB; CFBD; FB; **Synonyms** 

FBI12; GBG; H2-Bf; PBF2

**Host Species** 

Biotinylated Anti-CFB antibody(DMC366); IgG1 **Description** 

Chimeric mAb

Delivery 2-3 weeks **Uniprot ID** P00751

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended **Dilutions** 

**Background** 

Flow Cyt 1:100

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes complement factor B; a component of the alternative pathway of complement activation. Factor B circulates in the blood as a single chain polypeptide. Upon

activation of the alternative pathway; it is cleaved by complement factor D yielding the noncatalytic chain Ba and the catalytic subunit Bb. The active subunit Bb is a serine protease which associates with C3b to form the alternative pathway C3 convertase. Bb is involved in the proliferation of preactivated B lymphocytes; while Ba inhibits their proliferation. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. This cluster includes several genes involved in regulation of the

immune reaction. Polymorphisms in this gene are associated with a reduced risk of age-related macular degeneration. The polyadenylation site of this gene is 421 bp from the 5' end of the gene

for complement component 2.

Usage Research use only

Conjugate Biotinylated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

> > Email: info@dimabio.com

Website: www.dimabio.com

actively scrutinizing all patent application to

ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

**DIMA Disclaimer** 

/+86-400-006-0995(China)