

PRODUCT INFORMATION

Clone ID 55A1 HER3 **Target**

ERBB3; FERLK; LCCS2; VSCN1; ErbB-3; c-erbB3; erbB3-S; MDA-BF-1; c-erbB-3; p180-ErbB3; p45-**Synonyms**

sErbB3; p85-sErbB3

Host Species

Biotinylated Anti-Her3 antibody(55A1), Rabbit **Description**

mAb

2-3 weeks Delivery P21860 **Uniprot ID** Rabbit IgG IgG type Monoclonal Clonality Reactivity Human

Applications IHC; Flow Cyt

Recommended

Storage & Shipping

Background

IHC 1:100; Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before Formulation & lyophilization. Please see Certificate of Analysis Reconstitution for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or

overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been

characterized. One isoform lacks the

intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized. [provided by RefSeq, Jul

Email: info@dimabio.com

Website: www.dimabio.com

20081

Usage Research use only

Conjugate Biotinylated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)



Cat. No. DME100580B



DIMA Disclaimer

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to JIMARIO CONTRIBUTIONI DE LA CONTRIBUTIONI DEL CONTRIBUTIONI DEL CONTRIBUTIONI DE LA CONTRIBUTIONI DE LA CONTRIBUTIONI DE LA CONTRIBUTIONI DE LA CONTRIBUTIONI DEL CONTRIBUTIONI DE LA CONTRIBUTIONI DE LA CONTRIBUTIONI DE LA CONTRIBUTIONI DEL CONTRIBUTIONI DE LA CONTRI ensure no IP infringement.

> Email: info@dimabio.com Website: www.dimabio.com



