

PRODUCT INFORMATION

Clone ID 4G2 IL1A **Target**

Synonyms IL-1 alpha;IL-1A;IL1;IL1-ALPHA;IL1F1

Host Species

Biotinylated Anti-IL1A antibody(4G2); IgG1 Description

Chimeric mAb

Delivery 2-3 weeks **Uniprot ID** P01583

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Flow Cyt 1/100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is a pleiotropic cytokine involved in various immune

responses, inflammatory processes, and hematopoiesis. This cytokine is produced by monocytes and macrophages as a proprotein, which is proteolytically processed and released in response to cell injury, and thus induces apportunity. This gape and eight other interloukin 1

Background

apoptosis. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. It has been suggested that the polymorphism of these genes is associated with rheumatoid arthritis and Alzheimer's disease.

> Email: info@dimabio.com Website: www.dimabio.com

[provided by RefSeq, Jul 2008]

Usage Research use only

