Cat. No. DMC100467B



## **PRODUCT INFORMATION**

Clone ID **DMC467 Target** TGFBR2

AAT3; FAA3; LDS1B; LDS2; LDS2B; MFS2; RIIC; **Synonyms** TAAD2; TBR-ii; TBRII; TGFbeta-RII; TGFR-2

**Host Species** 

Biotinylated Anti-TGFBR2 antibody(DMC467); IgG1 Chimeric mAb **Description** 

Delivery 2-3 weeks **Uniprot ID** P37173

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a transmembrane protein that has a protein kinase

domain; forms a heterodimeric complex with TGF-beta receptor type-1; and binds TGF-beta. This receptor:ligand complex phosphorylates proteins; which then enter the nucleus and regulate the transcription of genes related to cell proliferation; cell cycle arrest; wound healing:

**Background** cell cycle arrest; wound healing;

immunosuppression; and tumorigenesis.

Mutations in this gene have been associated with Marfan Syndrome; Loeys-Deitz Aortic Aneurysm Syndrome; and the development of various types of tumors. Alternatively spliced transcript variants encoding different isoforms have been

> Email: info@dimabio.com Website: www.dimabio.com

characterized. [provided by RefSeq; Aug 2017]

**Usage** Research use only

Conjugate Biotinylated

