

## PRODUCT INFORMATION

<b>Target</b>	CD3D and CD3E
<b>Synonyms</b>	CD3-DELTA; IMD19; T3D and IMD18; T3E; TCRE
<b>Description</b>	Recombinant Cynomolgus CD3D Protein with C-terminal 10×His tag and Cynomolgus CD3E Protein with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q95LI8 and Q95LI5
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-10×His tag and C-Human Fc tag
<b>Molecular Characterization</b>	CD3D(Phe22-Ala105) 10×His tag and CD3E(Gln22-Asp117) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 10.9 and 37.0 kDa after removal of the signal peptide.
<b>Purity</b>	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	T-cell surface glycoprotein CD3 delta and CD3 epsilon chain, also known as CD3D and CD3E or CD3D and CD3E respectively, are single-pass type I membrane proteins. CD3D, together with CD3- epsilon(CD3E) , CD3-gamma and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.
<b>Usage</b>	Research use only






Figure 1. Cynomolgus CD3D and CD3E Heterodimer Protein, His Tag and hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

