

PRODUCT INFORMATION

MUC1 **Target**

Synonyms

ADMCKD, ADMCKD1, CA 15-3, CD227, EMA, H23AG, KL-6, MAM6, MCD, MCKD, MCKD1, MUC-1, MUC-1/SEC, MUC-1/X, MUC1/ZD, PEM, PEMT, PUM Recombinant Cynomolgus MUC1(380-500) protein

Description with C-terminal mouse Fc tag

Delivery

Uniprot ID XP_005541632.2

HEK293 Expression Host

Tag C-Mouse Fc tag

Molecular

Storage & Shipping

Background

MUC1(Leu380-Val500) mFc(Pro99-Lys330) Characterization

The protein has a predicted molecular mass of **Molecular Weight**

39.9 kDa after removal of the signal peptide. The apparent molecular mass of cMUC1(380-500)-mFc is approximately 35-55 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue **Purity**

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before Formulation & Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient témperature.

This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role

in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues. including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex.


The N-terminal alpha subunit functions in celladhesion and the C-terminal beta subunit is

involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results

> Email: info@dimabio.com Website: www.dimabio.com

in multiple transcript variants.

Usage Research use only Conjugate Unconjugated








Figure 1. Cynomolgus MUC1(380-500) Protein, mFc Tag on SDS-PAGE under reducing condition.

