

PRODUCT INFORMATION

Target	FZD10
Description	Monoclonal Cell Line Derived from K562 Cells, Engineered for Stable Expression of Human FZD10 Using Lentiviral Technology
Host Cells	K562
Uniprot ID	Q9ULW2
Applications	FACS Data
Growth media	RPMI-1640+10% FBS+1% P.S+1% Gln+2 ug/mL Puromycin
Package	5E6 Cells/mL
Suggested Control	SKU: BME100185
Warranty and Disclaimer	1. Please inspect cells upon receipt and report any issues promptly. 2. We offer one-time replacements for issues reported within a week of receipt. 3. User-induced issues are not eligible for free replacements. 4. We do not accept liability for damages resulting from cell use, storage, or loss. 5. Feedback received more than one month after receipt will not be processed.
Storage & Shipping	Cells are shipped using dry ice and require liquid nitrogen storage for long term preservation.
Synonyms	CD350; FZ-10; Fz10; FzE7; hFz10
Background	This gene is a member of the frizzled gene family. Members of this family encode 7-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. Using array analysis; expression of this intronless gene is significantly up-regulated in two cases of primary colon cancer. [provided by RefSeq; Jul 2008]
Usage	For research use only.



Hu_FZD10 K562 Cell Line




Figure 1. Flow cytometry analysis of human FZD10 overexpression using Hu_FZD10 K562 Cell Line (Cat. No. CEL100053) and Anti-FZD10(tabituximab biosimilar) mAb (Cat. No. BME100185)

DIMABIO CONFIDENTIAL

