


PRODUCT INFORMATION

Тад	C-Flag Tag
Target	ADGRE1
Synonyms	EMR1; TM7LN3
Description	Human ADGRE1 full length protein-synthetic
Delivery	3-4 weeks
Uniprot ID	Q14246
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length ADGRE1 protein has a MW of 97.5 kDa
Formulation & Reconstitution	
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes apolipoprotein A-I, which is the major protein component of high density lipoprotein (HDL) in plasma. The encoded preproprotein is proteolytically processed to generate the mature protein, which promotes cholesterol efflux from tissues to the liver for excretion, and is a cofactor for lecithin cholesterolacyltransferase (LCAT), an enzyme responsible for the formation of most plasma cholesteryl esters. This gene is closely linked with two other apolipoprotein genes on chromosome 11. Defects in this gene are associated with HDL deficiencies, including Tangier disease, and with systemic non-neuropathic amyloidosis. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein. [provided by RefSeq, Dec 2015]
Usage	Research use only
Conjugate	Unconjugated










Figure1. Elisa plates were pre-coated with Flag Tag ADGRE1-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ADGRE1-Nanodisc is 3.996ng/ml.




Figure2. Human ADGRE1-Nanodisc, Flag Tag on SDS-PAGE

Email: info@dimabio.com Website: www.dimabio.com

