

PRODUCT INFORMATION

| | |
|---|--|
| Target | ADGRE5 |
| Synonyms | CD97; TM7LN1 |
| Description | Human ADGRE5 full length protein-synthetic nanodisc |
| Delivery | In Stock |
| Uniprot ID | P48960 |
| Expression Host | HEK293 |
| Protein Families | Druggable Genome, GPCR, Transmembrane |
| Protein Pathways | N/A |
| Molecular Weight | The human full length ADGRE5 protein has a MW of 90.5 kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | A member of the EGF-TM7 subfamily of adhesion G protein-coupled receptors, which mediate cell-cell interactions. These proteins are cleaved by self-catalytic proteolysis into a large extracellular subunit and seven-span transmembrane subunit, which associate at the cell surface as a receptor complex. The encoded protein may play a role in cell adhesion as well as leukocyte recruitment, activation and migration, and contains multiple extracellular EGF-like repeats which mediate binding to chondroitin sulfate and the cell surface complement regulatory protein CD55. Expression of this gene may play a role in the progression of several types of cancer. Alternatively spliced transcript variants encoding multiple isoforms with 3 to 5 EGF-like repeats have been observed for this gene. This gene is found in a cluster with other EGF-TM7 genes on the short arm of chromosome 19. |
| Usage | Research use only |



ELISA assay to evaluate ADGRE5-Nanodisc 0.2 μ g Human ADGRE5-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag ADGRE5-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ADGRE5-Nanodisc is 7.327ng/ml.

kDa M R



Figure2. Human ADGRE5-Nanodisc, Flag Tag on SDS-PAGE

