

**PRODUCT INFORMATION**

<b>Target</b>	ADGRG1
<b>Synonyms</b>	BFPP; BPPR; GPR56; TM7LN4; TM7XN1
<b>Description</b>	Human ADGRG1 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9Y653
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, GPCR, Transmembrane
<b>Protein Pathways</b>	N.A.
<b>Molecular Weight</b>	The human full length ADGRG1 protein has a MW of 77.7 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C(Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	A member of the G protein-coupled receptor family and regulates brain cortical patterning. The encoded protein binds specifically to transglutaminase 2, a component of tissue and tumor stroma implicated as an inhibitor of tumor progression. Mutations in this gene are associated with a brain malformation known as bilateral frontoparietal polymicrogyria.
<b>Usage</b>	Research use only



### ELISA assay to evaluate ADGRG1-Nanodisc 0.2 $\mu$ g Human ADGRG1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag ADGRG1-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ADGRG1-Nanodisc is 1.419ng/ml.



Figure2. Human ADGRG1-Nanodisc, Flag Tag on SDS-PAGE

