

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	AGRB1
Synonyms	BAI1, GDAIF
Description	Human AGRB1 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	O14514
Expression Host	HEK293
Protein Families	Transmembrane,Druggable Genome,
Protein Pathways	Angiogenesis,Cancer,P53,Apoptosis & Cell Cycle,p53 Signaling Pathway,
Molecular Weight	The human full length AGRB1 protein has a MW of 173.5kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Angiogenesis is controlled by a local balance between stimulators and inhibitors of new vessel growth and is suppressed under normal physiologic conditions. Angiogenesis has been shown to be essential for growth and metastasis of solid tumors. In order to obtain blood supply for their growth, tumor cells are potently angiogenic and attract new vessels as results of increased secretion of inducers and decreased production of endogenous negative regulators. BAI1 contains at least one 'functional' p53-binding site within an intron, and its expression has been shown to be induced by wildtype p53. There are two other brain-specific angiogenesis inhibitor genes, designated BAI2 and BAI3 which along with BAI1 have similar tissue specificities and structures, however only BAI1 is transcriptionally regulated by p53. BAI1 is postulated to be a member of the secretin receptor family, an inhibitor of angiogenesis and a growth suppressor of glioblastomas [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

