

PRODUCT INFORMATION

Target	ANG1
Synonyms	AGP1;AGPT;ANG1
Description	Recombinant human ANG1 protein with C-terminal human Fc tag
Delivery	Under development
Uniprot ID	O76062
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	ANG1(Tyr378-Tyr418) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 40.04 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a secreted glycoprotein that belongs to the angiotensin family. Members of this family play important roles in vascular development and angiogenesis. All angiotensins bind with similar affinity to an endothelial cell-specific tyrosine-protein kinase receptor. The protein encoded by this gene is a secreted glycoprotein that activates the receptor by inducing its tyrosine phosphorylation. It plays a critical role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme and inhibits endothelial permeability. The protein also contributes to blood vessel maturation and stability, and may be involved in early development of the heart. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Sep 2015]
Usage	Research use only
Conjugate	Unconjugated

