

PRODUCT INFORMATION

| Тад | C-Flag Tag |
|---------------------------------|--|
| Target | ASIC4 |
| Synonyms | ACCN4, BNAC4 |
| Description | Human ASIC4 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q96FT7 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Other |
| Protein Pathways | N/A |
| Molecular Weight | The human full length ASIC4 protein has a MW of |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene belongs to the superfamily of acid- sensing ion channels, which are proton-gated, amiloride-sensitive sodium channels. These channels have been implicated in synaptic transmission, pain perception as well as mechanoperception. This gene is predominantly expressed in the pituitary gland, and was considered a candidate for paroxysmal dystonic choreoathetosis (PDC), a movement disorder, however, no correlation was found between mutations in this gene and PDC. [provided by RefSeq, Feb 2012] |
| Usage | Research use only |
| Conjugate | Unconjugated |
| | |

Email: info@dimabio.com Website: www.dimabio.com

