

PRODUCT INFORMATION

| Тад | C-Flag Tag |
|---------------------------------|---|
| Target | CAC1 |
| Synonyms | Cav3.3, ca(v)3.3 |
| Description | Human CAC1 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q9P0X4 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Calcium |
| Protein Pathways | N/A |
| Molecular Weight | The human full length CAC1 protein has a MW of 245.1kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes the pore-forming alpha subunit of a voltage gated calcium channel. The encoded protein is a member of a subfamily of calcium channels referred to as is a low voltage- activated, T-type, calcium channel. The channel encoded by this protein is characterized by a slower activation and inactivation compared to other T-type calcium channels. This protein may be involved in calcium signaling in neurons. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Oct 2011] |
| Usage | Research use only |
| Conjugate | Unconjugated |
| | |

Email: info@dimabio.com Website: www.dimabio.com

