

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Tag</b>                              | C-Flag&Strep Tag  |
| <b>Target</b>                           | CAC15   |
| <b>Synonyms</b>                         | CACNL1A3, CCHL1A3, Cav1.1, HOKPP, HOKPP1, MHS5, TTPP1, hypoPP   |
| <b>Description</b>                      | Human CAC15-Strep full length protein-synthetic nanodisc  |
| <b>Delivery</b>                         | 6~8weeks  |
| <b>Uniprot ID</b>                       | Q13698  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Protein Families</b>                 | Ion Channels: Calcium   |
| <b>Protein Pathways</b>                 | N/A   |
| <b>Molecular Weight</b>                 | The human full length CAC15-Strep protein has a MW of 212.4 kDa<br>Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| <b>Formulation &amp; Reconstitution</b> | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Storage &amp; Shipping</b>           |   |
| <b>Background</b>                       | This gene encodes one of the five subunits of the slowly inactivating L-type voltage-dependent calcium channel in skeletal muscle cells. Mutations in this gene have been associated with hypokalemic periodic paralysis, thyrotoxic periodic paralysis and malignant hyperthermia susceptibility. [provided by RefSeq, Jul 2008]   |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |

