

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CCG2
Synonyms	MRD10
Description	Human CCG2 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9Y698
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length CCG2 protein has a MW of 36kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a type I transmembrane AMPA receptor regulatory protein (TARP). TARPs regulate both trafficking and channel gating of the AMPA receptors. The AMPA subtype of ionotropic glutamate receptors are ligand gated ion channels that are typically activated by glutamate released from presynaptic neuron terminals and mediate fast neurotransmission in excitatory synapses. TARPs thus play an important role in synaptic plasticity, learning and memory. Mutations in this gene cause an autosomal dominant form of cognitive disability. [provided by RefSeq, Jul 2017]
Usage	Research use only
Conjugate	Unconjugated

