

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

CCR10 **Target** GPR2 **Synonyms**

Formulation &

Reconstitution

Storage & Shipping

Background

Human CCR10-Strep full length protein-synthetic **Description**

nanodisc **Delivery** 6~8weeks **Uniprot ID** P46092 **Expression Host HEK293**

Protein Families Druggable Genome, GPCR, Transmembrane

Chemokine signaling pathway, Cytokine-cytokine **Protein Pathways**

receptor interaction

The human full length CCR10-Strep protein has a **Molecular Weight**

MW of 38.4 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Lyophilized from PBS. Normally 5% - 8% trehalose is added as

protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in

subsequent experiments. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

Chemokines are a group of small (approximately 8 to 14 kD), mostly basic, structurally related molecules that regulate cell trafficking of various types of leukocytes through interactions with a subset of 7-transmembrane, G protein-coupled receptors. Chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in

angiogenesis or angiostasis. Chemokines are

divided into 2 major subfamilies, CXC and CC, based on the arrangement of the first 2 of the 4 conserved cysteine residues; the 2 cysteines are separated by a single amino acid in CXC

chemokines and are adjacent in CC chemokines. CCR10 is the receptor for CCL27 (SCYA27; MIM 604833); CCR10-CCL27 interactions are involved

> Email: info@dimabio.com Website: www.dimabio.com

in T cell-mediated skin inflammation.

Usage Research use only Conjugate Unconjugated



