

## **PRODUCT INFORMATION**

Тад	C-Flag&Strep Tag
Target	CCR5
Synonyms	CC-CKR-5; CCCKR5; CCR-5; CD195; CKR-5; CKR5; CMKBR5; IDDM22
Description	Human CCR5-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P51681
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, ES Cell Differentiation/IPS, GPCR, Transmembrane
Protein Pathways	Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Endocytosis
Molecular Weight	The human full length CCR5-Strep protein has a MW of 40.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. This protein is expressed by T cells and macrophages, and is known to be an important co-receptor for macrophage-tropic virus, including HIV, to enter host cells. Defective alleles of this gene have been associated with the HIV infection resistance. The ligands of this receptor include monocyte chemoattractant protein 2 (MCP-2), macrophage inflammatory protein 1 alpha (MIP-1 alpha), macrophage inflammatory protein 1 beta (MIP-1 beta) and regulated on activation normal T expressed and secreted protein (RANTES). Expression of this gene was also detected in a promyeloblastic cell line, suggesting that this protein may play a role in granulocyte lineage proliferation and differentiation. This gene is located at the chemokine receptor gene cluster region. An allelic polymorphism in this gene results in both functional and non-functional alleles; the reference genome represents the functional allele.
Usage	Research use only
Conjugate	Unconjugated

Email: info@dimabio.com Website: www.dimabio.com

