

PRODUCT INFORMATION

Target	CD151
Synonyms	GP27; MER2; PETA-3; RAPH; SFA1; TSPAN24
Description	Human CD151 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P48509
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length CD151 protein has a MW of 28.3 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins and other transmembrane 4 superfamily proteins. It is involved in cellular processes including cell adhesion and may regulate integrin trafficking and/or function. This protein enhances cell motility, invasion and metastasis of cancer cells. Multiple alternatively spliced transcript variants that encode the same protein have been described for this gene.
Usage	Research use only



ELISA assay to evaluate CD151-Nanodisc
0.2µg Human CD151-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CD151-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CD151-Nanodisc is 4.446ng/ml.



Figure2. Human CD151-Nanodisc, Flag Tag on SDS-PAGE

