

PRODUCT INFORMATION

Target	CLDN18.2
Synonyms	SFTA5; SFTPJ
Description	Human CLDN18.2 full length protein-MNP
Delivery	In Stock
Uniprot ID	P56856
Expression Host	HEK293
Protein Families	Transmembrane
Protein Pathways	Cell adhesion molecules (CAMs), Leukocyte transendothelial migration, Tight junction
Molecular Weight	The human full length CLDN18.2 Protein has a MW of 27.5 kDa
Formulation & Reconstitution	Lyophilized from PBS. Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A member of the claudin family. Claudins are integral membrane proteins and components of tight junction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. This gene is upregulated in patients with ulcerative colitis and highly overexpressed in infiltrating ductal adenocarcinomas. PKC/MAPK/AP-1 (protein kinase C/mitogen-activated protein kinase/activator protein-1) dependent pathway regulates the expression of this gene in gastric cells. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jun 2010]
Usage	Research use only
Conjugate	Unconjugated



**ELISA assay to evaluate CLDN18.2-MNP**  
0.5µg Human CLDN18.2-MNP per well




Figure1. Elisa plates were pre-coated with 0.5µg/per well purified human CLDN18.2 full length membrane nanoparticles. Serial diluted anti-CLDN18.2 monoclonal antibody (BME100075) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-CLDN18.2 monoclonal antibody binding with CLDN18.2 full length membrane nanoparticles is 16.39ng/ml.

