

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag Tag |
| Target | CLIC5 |
| Synonyms | DFNB102, DFNB103, MST130, MSTP130 |
| Description | Human CLIC5 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q9NZA1 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Other |
| Protein Pathways | N/A |
| Molecular Weight | The human full length CLIC5 protein has a MW of 46.5kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a member of the chloride intracellular channel (CLIC) family of chloride ion channels. The encoded protein associates with actin-based cytoskeletal structures and may play a role in multiple processes including hair cell stereocilia formation, myoblast proliferation and glomerular podocyte and endothelial cell maintenance. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2011] |
| Usage | Research use only |
| Conjugate | Unconjugated |

