

PRODUCT INFORMATION

Target	CMTM6
Synonyms	CKLFSF6; PRO2219
Description	Human CMTM6 full length protein-synthetic nanodisc
Delivery	3-4 weeks
Uniprot ID	Q9NX76
Expression Host	HEK293
Protein Families	Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length CMTM6 protein has a MW of 20.4 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene belongs to the chemokine-like factor gene superfamily, a novel family that is similar to the chemokine and transmembrane 4 superfamilies. This gene is one of several chemokine-like factor genes located in a cluster on chromosome 3. This gene is widely expressed in many tissues, but the exact function of the encoded protein is unknown. [provided by RefSeq, Jul 2008]
Usage	Research use only



ELISA assay to evaluate CMTM6-Nanodisc 0.2 μ g Human CMTM6-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CMTM6-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CMTM6-Nanodisc is 4.393ng/ml.



Figure2. Human CMTM6-Nanodisc, Flag Tag on SDS-PAGE

