

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Tag</b>                              | C-Flag Tag  |
| <b>Target</b>                           | CNGA2   |
| <b>Synonyms</b>                         | CNCA, CNCA1, CNG2, OCNC1, OCNCALPHA, OCNCa  |
| <b>Description</b>                      | Human CNGA2 full length protein-synthetic nanodisc  |
| <b>Delivery</b>                         | 6~8weeks  |
| <b>Uniprot ID</b>                       | Q16280  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Protein Families</b>                 | Ion Channels: Cyclic nucleotide gated   |
| <b>Protein Pathways</b>                 | N/A   |
| <b>Molecular Weight</b>                 | The human full length CNGA2 protein has a MW of 76kDa   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for   |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| <b>Background</b>                       | The protein encoded by this gene represents the alpha subunit of a cyclic nucleotide-gated olfactory channel. The encoded protein contains a carboxy-terminal leucine zipper that mediates channel formation. [provided by RefSeq, Jan 2010]        |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |

