

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CTSR2
Synonyms	N/A
Description	Human CTSR2 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q96P56
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length CTSR2 protein has a MW of 62kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of a family of cation channel proteins that localize to the flagellum of spermatozoa. Defects at this locus causes male infertility. Alternatively spliced transcript variants have been observed at this locus. Readthrough transcription originates upstream of this locus in diphosphoinositol pentakisphosphate kinase 1 pseudogene 1 and is represented by GeneID:110006325. Related pseudogenes are found next to this locus on chromosome 15 and on chromosome 5. [provided by RefSeq, Mar 2017]
Usage	Research use only
Conjugate	Unconjugated

